

THE SOLUTION PHYSICIANS TRUST AND PATIENTS COUNT ON



Long-term clinical trial data

U.S. IDE Clinical Trial 5-year data
63 patients* from primary enrollment

100%

Patency — External iliac artery

95.1%

Patency — Internal iliac artery

0%

Buttock claudication†

0%

New onset erectile dysfunction

ZERO

Type I/III endoleaks
Migrations

98.3%

Freedom from CIAA†
enlargement (> 5 mm)

95.2%

Freedom from IBE-related
reintervention

Core Lab reported assessment for patency, endoleak, migration and CIAA enlargement. Denominator is number of subjects evaluated for primary effectiveness endpoint result with an evaluable result.

* Sixty-three subjects with device implanted in initial cohort. Thirty-six patients have completed 5-year follow-up.

† On the side treated with the IBE.

Together, improving life



THE SVS CLINICAL PRACTICE GUIDELINES FOR ABDOMINAL AORTIC ANEURYSM¹ (AAA) INCLUDE RECOMMENDATIONS PERTAINING TO HYPOGASTRIC PRESERVATION DURING EVAR TREATMENT.

These recommendations include:

Preservation of flow to at least one internal iliac artery

1 (Strong)

Level of recommendation

A (High)

Quality of evidence

Using a Food and Drug Administration (FDA) approved branch endograft in anatomically suitable patients to maintain perfusion to at least one internal iliac artery.

1 (Strong)

Level of recommendation

A (High)

Quality of evidence

PRESERVE FLOW. ADVANCE CARE.

- **Preservation matters:** Recommended treatment^{1,2} to sustain quality of life
- **Advances repair:** All-In-One System with proven outcomes
- **Performs as promised:** High patency³, conformability and durability
- **Complete confidence:** More than 20 years of aortic device experience



Learn more at goremedical.com/IBE

Cover image: 5-year follow-up; first clinical use. Image courtesy of Brian Peterson, MD., St. Anthony's Medical Center; St. Louis, Missouri.

Additional clinical trial data available in *Instructions for Use*. Additional clinical study information available.³

1. Chaikof EL, Dalman RL, Eskandari MK, et al. The Society for Vascular Surgery practice guidelines on the care of patients with an abdominal aortic aneurysm. *Journal of Vascular Surgery* 2018;67(1):2-77.e2.
2. Moll F.L., Powell J.T., Fraedrich G., et al; European Society for Vascular Surgery. Management of abdominal aortic aneurysms clinical practice guidelines of the European Society for Vascular Surgery. *European Journal of Vascular & Endovascular Surgery* 2011;41(Supplement 1):S1-S58.
3. Schneider DB, Matsumura JS, Lee JT, Peterson BG, Chaer RA, Oderich GS. Final 5-year results of the United States prospective, multicenter study of endovascular repair of iliac aneurysms using the Gore Iliac Branch Endoprosthesis. Presented at the Vascular Annual Meeting (VAM21); August 18-21; San Diego, CA. Abstract RS13.

 Consult Instructions for Use
eifu.goremedical.com

Refer to *Instructions for Use* at eifu.goremedical.com for a complete description of all applicable indications, warnings, precautions and contraindications for the markets where this product is available. R_{only}

Products listed may not be available in all markets.

GORE, *Together, improving life*, EXCLUDER and designs are trademarks of W. L. Gore & Associates.
© 2021 W. L. Gore & Associates, Inc. 21263094-EN OCTOBER 2021

W. L. Gore & Associates, Inc.
goremedical.com

Asia Pacific +65 67332882 Australia / New Zealand 1800 680 424 Europe 00800 6334 4673
United States Flagstaff, AZ 86004 800 437 8181 928 779 2771

