## **Designed For Conformability**


Building Confidence in the Treatment of Type B Dissection




## Worldwide data supports performance in the treatment of all Type B dissections

Gore GREAT Registry 1 provides worldwide experience and outcomes in real-world dissection use.

## **GREAT Dissection Experience**







Adverse Event	GREAT	Data <sup>3</sup>

Dissection-Related Survival	96.4 %
Procedural Survival	100 %
RTAD	0.6 %
Stroke	1.2 %
Type IA Endoleak	0 %
Device Compressions	0 %
Paraparesis / Paralysis	1.2 %
Conversion	0.6 %



- 2. The GORE® TAG® Thoracic Endoprosthesis is not indicated for the treatment of Zone 0 and Zone 1.
- 3. Includes all Type B dissections, events occurring within 12 months post-procedure.



Images courtesy of Nimesh Desai, MD

INDICATIONS FOR USE IN THE US: The GORE® TAG® Thoracic Endoprosthesis is intended for endovascular repair of all lesions of the descending thoracic aorta, including: Isolated lesions in patients who have appropriate anatomy, including: adequate iliac / femoral access, aortic inner diameter in the range of  $16-42\,\mathrm{mm}_* \ge 20\,\mathrm{mm}$  non-aneurysmal aorta proximal and distal to the lesion; Type B dissections in patients who have appropriate anatomy, including: adequate iliac / femoral access,  $\ge 20\,\mathrm{mm}$  landing zone proximal to the primary entry tear; proximal extent of the landing zone must not be dissected, diameter at proximal extent of proximal landing zone in the range of  $16-42\,\mathrm{mm}$ . CONTRAINDICATIONS: Patients with known sensitivities or allergies to the device materials; patients who have a condition that threatens to infect the graft. Refer to Instructions for Use at goremedical.com for a complete description of all warnings, precautions, and adverse events. INDICATIONS FOR USE UNDER CE MARK: The GORE® TAG® Thoracic Endoprosthesis is indicated for endovascular repair of the descending thoracic aorta. CONTRAINDICATIONS: Patients with known sensitivities or allergies to the device materials; patients with a systemic infection who may be at increased risk of endovascular graft infection. Refer to Instructions for Use at goremedical.com for a complete description of all warnings, precautions, and adverse events.  $\frac{R}{N}$ 

