




The solution physicians trust and patients count on

2-year data now available



The recently released SVS Clinical Practice Guidelines for Abdominal Aortic Aneurysm* (AAA) include updated recommendations pertaining to hypogastric preservation during EVAR treatment. These recommendations include:

Preservation of flow to at least one internal iliac artery

- Level of recommendation: 1 (Strong)
- Quality of evidence: A (High)

NEW! Using a Food and Drug Administration (FDA) approved branch endograft in anatomically suitable patients to maintain perfusion to at least one internal iliac artery.

- Level of recommendation: 1 (Strong)
- Quality of evidence: A (High)

Learn more at goremedical.com/IBE

* Chaikof EL, Dalman RL, Eskandari MK, *et al.* The Society for Vascular Surgery practice guidelines on the care of patients with an abdominal aortic aneurysm. *Journal of Vascular Surgery* 2018;67(1):2-77.e2.



GORE® EXCLUDER® Iliac Branch Endoprosthesis — Mid-term clinical trial data

U.S. IDE Clinical Trial now has 2-year follow-up data for all patients from primary enrollment (n = 63)

Key research outcomes are:

2-year data

Patency — External iliac artery	100%
Patency — Internal iliac artery	95.1%
Freedom from reintervention	93.7%
Buttock claudication	0%
New onset erectile dysfunction	0%
Freedom from CIAA enlargement (> 5 mm)	98.3%

152 minutes

Procedure time



40 minutes

Fluoro time



114 ml

Contrast used



95.2%


Technical success*



*Defined as successful implantation with lack of endoleaks

Preserve flow. Advance care.

- **Preservation matters:** Recommended treatment^{1,2} to sustain quality of life
- **Advances repair:** All-In-One System with proven outcomes
- **Performs as promised:** High patency³, conformability, and durability
- **Complete confidence:** More than 20 years of aortic device experience



Cover image: 3-year follow-up; first clinical use. Image courtesy of Brian Peterson, MD., St. Anthony's Medical Center; St. Louis, Missouri.

Additional clinical trial data available in *Instructions for Use*.
Additional clinical study information available.³

1. Chaikof EL, Dalman RL, Eskandari MK, et al. The Society for Vascular Surgery practice guidelines on the care of patients with an abdominal aortic aneurysm. *Journal of Vascular Surgery* 2018; 67(1):2-77.e2.
2. Moll F.L., Powell J.T., Fraedrich G., et al; European Society for Vascular Surgery. Management of abdominal aortic aneurysms clinical practice guidelines of the European Society for Vascular Surgery. *European Journal of Vascular & Endovascular Surgery* 2011;41(Supplement 1):S1-S58.
3. Schneider D. B. One-year U.S. results of GORE® EXCLUDER® Iliac Branched Endograft: Advantages and limitations. Presented at the 42nd Annual Symposium on Vascular and Endovascular Issues, Techniques, *Horizons* (VEITHsymposium); November 17-21, 2015; New York, NY.

Products listed may not be available in all markets.

GORE®, EXCLUDER®, and designs are trademarks of W. L. Gore & Associates.
© 2018 W. L. Gore & Associates, Inc. AX0410-EN4 MAY 2018



W. L. GORE & ASSOCIATES, INC.
Flagstaff, AZ 86004

+65.67332882 (Asia Pacific)
00800.6334.4673 (Europe)
800.437.8181 (United States)
928.779.2771 (United States)

goremedical.com