

# The solution physicians trust and patients count on



## GORE® EXCLUDER® Iliac Branch Endoprosthesis — New Mid-term clinical trial data

U.S. IDE Clinical Trial now through 3-year follow-up data  
for all patients from primary enrollment (n = 63)

|  |              |
|--|--------------|
| Patency — External iliac artery        | <b>100%</b>  |
| Patency — Internal iliac artery        | <b>95.1%</b> |
| Freedom from reintervention            | <b>93.7%</b> |
| Buttock claudication                   | <b>0%</b>    |
| New onset erectile dysfunction         | <b>0%</b>    |
| Freedom from CIAA enlargement (> 5 mm) | <b>98.3%</b> |

**152 minutes**

Procedure time



**40 minutes**

Fluoro time



**114 ml**

Contrast used



**95.2%**

Technical success\*



\* Defined as successful implantation  
with lack of endoleaks

The SVS Clinical Practice Guidelines for Abdominal Aortic Aneurysm\* (AAA) include recommendations pertaining to hypogastric preservation during EVAR treatment.

These recommendations include:

### Preservation of flow to at least one internal iliac artery

**1 (Strong)**

Level of recommendation

**A (High)**

Quality of evidence

**Using a Food and Drug Administration (FDA) approved branch endograft in anatomically suitable patients to maintain perfusion to at least one internal iliac artery.**

**1 (Strong)**


Level of recommendation

**A (High)**

Quality of evidence

## Preserve flow. Advance care.

- **Preservation matters:** Recommended treatment<sup>1,2</sup> to sustain quality of life
- **Advances repair:** All-In-One System with proven outcomes
- **Performs as promised:** High patency<sup>3</sup>, conformability and durability
- **Complete confidence:** More than 20 years of aortic device experience



### Learn more at [goremedical.com/IBE](http://goremedical.com/IBE)

\* Chaikof EL, Dalman RL, Eskandari MK, *et al.* The Society for Vascular Surgery practice guidelines on the care of patients with an abdominal aortic aneurysm. *Journal of Vascular Surgery* 2018;67(1):2-77.e2.

Cover image: 3-year follow-up; first clinical use. Image courtesy of Brian Peterson, MD., St. Anthony's Medical Center; St. Louis, Missouri.

Additional clinical trial data available in *Instructions for Use*.  
Additional clinical study information available.<sup>3</sup>

1. Chaikof EL, Dalman RL, Eskandari MK, *et al.* The Society for Vascular Surgery practice guidelines on the care of patients with an abdominal aortic aneurysm. *Journal of Vascular Surgery* 2018; 67(1):2-77.e2.
2. Moll F.L., Powell J.T., Fraedrich G., *et al.*; European Society for Vascular Surgery. Management of abdominal aortic aneurysms clinical practice guidelines of the European Society for Vascular Surgery. *European Journal of Vascular & Endovascular Surgery* 2011;41(Supplement 1):S1-S58.
3. Schneider D. B. One-year U.S. results of GORE® EXCLUDER® Iliac Branched Endograft: Advantages and limitations. Presented at the 42nd Annual Symposium on Vascular and Endovascular Issues, Techniques, *Horizons* (VEITHsymposium); November 17-21, 2015; New York, NY.

Products listed may not be available in all markets.

GORE, EXCLUDER, and designs are trademarks of W. L. Gore & Associates.

© 2018, 2019 W. L. Gore & Associates, Inc. AX0410-EN5 SEPTEMBER 2019



**W. L. GORE & ASSOCIATES, INC.**  
Flagstaff, AZ 86004

+65.67332882 (Asia Pacific)  
00800.6334.4673 (Europe)  
800.437.8181 (United States)  
928.779.2771 (United States)

[goremedical.com](http://goremedical.com)